

Study on the Nutritional Status of HIV Infected Adults

I.Devika P.hD Scholar, Dr.Thahira Banu.A

Department of Home Science, Gandhigram Rural Institute. Dindigul district.

Abstract:

The study was undertaken to assess the nutritional status of HIV subjects in Dindigul District, Tamilnadu, India. The nutritional status was assessed in 100 subjects. The results revealed that fifty three percent of the subjects had low level of CD₄ cell count, hemoglobin and dietary intake.

key words – HIV, Nutritional status, CD₄ count

INTRODUCTION:

The world's biggest challenges today emerges from four simple letters AIDS which stands for Acquired Immuno Deficiency Syndrome. The rate at which humanity falls prey to this plaque needs serious introspection. In India one percent of the total population is infected with HIV are through unethical sex and drug addiction being are the two common causes of AIDS. Nutrition plays a critical role in comprehensive care and support for people living with HIV. Nutritional interventions can help manage symptoms, promote response to medical treatment, slow progression of the disease and increase the quality of living. Dindigul district alone had over 8000 persons living with HIV but only 3411 subjects had registered with Anti- retroviral therapy centre of which 1,687 subjects got regular treatment (The Hindu, 2010). Hence this investigation was undertaken with an objective to assess the nutritional status of the HIV/infected subjects registered at ICTC centre in Dindigul District.

MATERIALS AND METHODS:

Selection of subjects:

A total of hundred HIV infected persons, in the age group of 20 to 55 years who were willing to participate in this study were selected using purposive sampling method from government hospital, Dindigul district; Tamil Nadu, India. All the selected hundred samples were assessed for their nutritional status. Among the selected subjects 56 were males and 44 were females.

Socioeconomic status:

The information regarding age, gender, educational qualifications, size of the family, type of the

family, income and occupation of the selected HIV subjects were collected using a well structured interview schedule.

Nutritional status

- Anthropometric measurement:* Based on the height and weight of the subjects the Body Mass Index (BMI) was calculated and the values were then compared with World Health Organization standards for under weight, normal and overweight. $BMI = \text{weight (kg)} / \text{height m}^2$. Height and weight are physical dimensions of the body that reflect the nutritional status. The height and weight of all the samples were recorded.
- Clinical assessment:* Clinical assessments were carried out with HIV subjects, using the formulated interview schedule. A thorough history and physical examinations with attention to symptoms manifested on the skin, nail, hair, stomach, respiratory symptoms and eyes and other symptoms were observed and recorded.
- Biochemical assessment:* Biochemical estimation was done by trained medical laboratory technician according to the protocol to be followed for testing HIV patient's blood. The immunity levels of the subjects were tested using the CD₄ count and blood hemoglobin.
- Dietary factors:* Data regarding the nature, quality and quantity of food consumed were collected using a 24 hour recall method. The raw equivalents was calculated and the nutrients namely energy, carbohydrate, protein, fat, iron, beta carotene, vitamin C and folic acid was computed by using the food consumption table from Nutritive Value of Indian Foods given by ICMR (Gopalan, 2007).

RESULTS AND DISCUSSION

Socio economic status:

Table 1 indicates that most of the subjects 57 percent were in their the age group of 30 to 39 years followed by 27, 8 and 8 percent of subjects in 40 to 49, 50 to 55 and 20 to 29 years respectively. Majority of 56 percent of subjects were male and only 44 percent of them were females.

Forty percent of the subjects had completed high school education and 17 subjects had only primary school education, five percent of the subjects had higher secondary education and only one subject was a graduate and 37

subjects were illiterates. The monthly income of majority of the subjects was Rs 1000 to 4000 per month and for 23 subjects it ranged from Rs 4000 to 7000 and for 4 subjects it was above 10,000 per month. Major occupation of the subjects was found to be coolies. In resource poor countries, many people live below poverty line and there is often no medical insurance or disability pension for people living with HIV (Katabira, 2002). Ninety four percent of the subjects were in nuclear family and 68 percent of the subjects had small family size.

Table 1

SOCIO ECONOMIC STATUS OF THE SELECTED

Particulars	Subjects N=100	Particulars	Subjects N=100
Age 20-29 30-39 40-49 50-55	8 57 27 8	Type of the family Nuclear joint	94 6
Gender Male Female	56 44	Income of the family 1000-4000 4000-5000 7000-1000 and above	73 23 4
Educational qualification Illiterate 1-5 Standard 6-10 Standard 11-12 Standard Under graduate	37 17 40 5 1		
Size of the family Small (1-3) Medium(4-5) Large (5 and above)	68 32 -	Occupation Agriculture Cooli Business	20 62 18

SUBJECTS

Nutritional status of the selected HIV subjects:

Body Mass Index:

It was found that 29 percent of male and 12 percent of female selected subjects were underweight, 24 percent of male and 28 percent of female subjects had normal Body Mass Index of 18.6 to 24.9 and six percent (3 male and 3female) of the subjects had a Body Mass Index of 25.0 to 29.9 and was in grade I obesity and one percent of female subject had grade II obesity. Obi *et al.* (2010) conducted a

study on nutritional status of HIV-positive individuals on free HAART treatment in a developing nation and found that there malnourished individuals among the HIV-positive group than in the control group ($P < 0.05$) it concluded that malnutrition is common among HIV-positive subjects in southeast Nigeria.

Clinical assessment:

Clinical assessment is an important practical method for assessing the nutritional status of the community. The clinical assessment reveals that 53percent (31 male and 22 female) subjects had distaste for food, 56 percent (34 male and 22 female) subjects had wasting of muscle, 41 percent (28 male and 13female) had some skin infections,31 percent (19 male and 12 female) of the subjects had tooth caries, 48 percent (31 male and 17 female) had early satiety, 52 percent (37 male and 15 female) had thin-sparse hair, 45 percent (36 male and 19 female) had pale conjunctiva, 49 percent (31 male and 18 females) had spoon shaped nails, 35 percent (25 male and 10 female) had discoloration of eyes, 38 percent (22 male and 16 female) had bleeding gums, 32 percent (20 male and 12 female) had skin ulcers and 22 percent (13 male and 09 female) of the subjects had nausea .

a)Biochemical assessment:

The Table 2 result shows that all the subjects had haemoglobin levels below the normal level of haemoglobin (12.9 g/dl).

Table 2

BIOCHEMICAL PARAMETERS OF SELECTED HIV INFECTED SUBJECTS

Biochemical parameters	Subjects n= 100		Reference value *	
	Male	Female	Male	Female
Haemoglobin Level			13.5- 17.91 g/dl	12.0 – 15.5 g/dl)
Below 6.9	02	01		
7 - 9.9	42	38		
10 -12.9	12	05		
CD₄ Cell count			(400- 1600 mm ³)	(400- 1600 mm ³)
Below 250	04	08		
251 -500	11	09		
501 – 1000	13	08		
1001 – 1500	09	07		
1501 and above	19	12		

*ICMR (2001)

a) Mean nutrient intake of the selected subjects:

The mean nutrient intake for the selected male subjects were found to be as follows, calories 3210 kcal, protein 42g, fat 29g, calcium 303 mg, iron 16 mg, beta carotene 819 µg, vitamin C 28 mg folic acid 32 mg, and 3 mg zinc. The mean nutrient intake has been compared with RDA, which reflects the percent adequacy of energy (11) and fat (44) . All the other nutrients were deficit in selected 100 subjects, other nutrients namely protein, calcium, iron, beta carotene, vitamin C, folic Acid and zinc by - 30, - 24 - 41, - 66, - 30, and - 68, and -67 percent respectively.

The mean nutrient intake for the selected female subjects were found to be as follows, calories 2874 kcal, protein 33.6g, fat 26.9g, calcium 382.4mg, iron 18.1 mg, beta carotene 873.6 µg, vitamin C 30.4mg , folic acid 30.9 mg and 3.6 mg zinc. The mean nutrient intake has been compared with RDA, which reflects the percent adequacy of energy 29.1 and fat 34.5 respectively. All the other nutrients were deficit in selected 100 subjects, other nutrients namely protein, calcium, iron, beta carotene, vitamin C, folic Acid, and zinc by - 32, - 4.5, - 9.6, - 63.6, - 24, and - 69.1, and – 44 respectively. The results clearly point out that all the subjects had a deficient intake of protein, calcium, iron, beta carotene, vitamin C, folic acid, and Zinc. HIV-infected individuals have higher rates of malnutrition, anaemia, and hypo albuminemia than socioeconomically matched HIV-uninfected individuals and despite similar caloric intake (Swaminathan *et al.* 2008).

CONCLUSION:

All the subjects had low level of hemoglobin and low level of nutrient intake except for energy and fat. There is need to give diet counseling to the subjects regarding nutritional status of the HIV infected subjects to reduce the opportunistic infections and improve their health status.

ACKNOWLEDGEMENT:

I wish to express my deep sense of gratitude and hearty thanks to my honorable Guide Mrs.A.Thahira Banu Assistant professor Department of Home Science,GRI and Heart full regards to my dear beloved parents, my family members for their love, concern and constant encouragement and support given throughout the study.

References :

- i) Gopalan,C.,Ramsastri,B.Vand alasubramaniS.C,(2007). *Nutritive value of Indian foods, NIN, ICMR:Hyderabad:78-81.*
- ii) Katabira, ET. (2002). *The promise and challenges of antiretroviral therapy in developing countries. Paper presented at the 19th conference on retroviruses and opportunistic infection. Symposium session 30.*
- iii) Obi, SN., Ifebunandu,NA., and Onyebuchi, AK. (2010). *Nutritional status of HIV-positive individuals on free HAART treatment in a developing nation. J Infect Dev Ctries. 2010 Nov 24;4(11):745-9.*
- iv) *Staff report (2010, May/6). Register with ARTcentres to avail support services. The Hindu.*
- v) Swaminathan, S., Padmapriyadarsini, C., and Sukumar, B. (2008). *Nutritional status of persons with HIV infection, persons with HIV infection and tuberculosis, and HIV-negative individuals from southern India. Clin Infect Dis;46:946-949.*